

# Minor II: CLL113 Numerical Methods in Chemical Engineering

Date: 10/10/16

Total Marks: 15

Please do not seek *any* clarification. Make suitable assumptions wherever necessary.

1. Determine all the values of  $a, b, c, d, e$  and  $f$  for which the following function is a cubic spline:

$$f(x) = \begin{cases} ax^2 + b(x-1)^3 & x \in [0, 1]; \\ cx^2 + d & x \in [1, 2]; \\ ex^2 + f(x-2)^3 & x \in [2, 3]. \end{cases} \quad [3]$$

2. Consider the approximation to the following integration:

$$\int_{-1}^1 f(x) dx \approx f(\alpha) + f(-\alpha).$$

For what value(s) of  $\alpha$ , if any, will this formula be *exact* for the function  $f(x)$  being all polynomials of the form  $a + bx + cx^3 + dx^4$ , where  $a, b, c$  and  $d$  are constants? [4]

3. For two non-linear algebraic equations:  $f_1(x, y) = 0$  and  $f_2(x, y) = 0$ , the programming function "*core*( $f_1, f_2, x, y$ )" returns the value of functions  $f_1$  and  $f_2$  for input values of  $x$  and  $y$ . Write an algorithm for Newton-Raphson technique to find the roots  $x$  and  $y$  starting with initial guess  $x_0$  and  $y_0$ . Use central difference formula to construct the Jacobian matrix. The convergence condition should be based on relative (or normalized) error. Show only the important steps. [3]

4. Consider the data points:

$x$	0	1	2	4	5	6
$f(x)$	1	14	15	5	6	19

Using the Newton-divided difference method, obtain coefficients of 7<sup>th</sup> degree polynomial  $a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7$  which passes through *all* the points and find  $f(3)$ . Find the derivative  $f'(1)$  using any 2<sup>nd</sup> order accurate formula. Give at least two reasons why this value of derivative is not in agreement with the  $f'(1)$  calculated from the fitted polynomial (exact value). [2+1.5]

5. Answer briefly:

2'

(a) For fitting function values at  $n$  data point where  $n$  is very large, explain why cubic spline is preferable over single interpolating polynomial of either Lagrange or Newton type? [1]

(b) Briefly explain why numerical differentiation is less accurate computationally as compared to numerical integration. [0.5]

( )